

DAFTAR PUSTAKA

- Barbosa, L., Berk, M., Vorster, M. (2003). *A Double – Blind randomized, placebo- controlled trial of augmentation with lamotrigine or placebo in patients episodes.* <http://www.ncbi.nlm.nih.gov/pubmed/12716240>
- Beck, J. S. (2011). *Cognitive behavior therapy: Basics and beyond.* Guilford press.
- Center for CBT. *Post traumatic stress disorder.* 31 Maret 2010. Diakses di : <http://www.centreforcbtcounselling.co.uk/ptsd.php>. (diakses tanggal 03 Mei 2020).
- David D, Szentagotai A, Lupu V, dkk. *Rational emotive behavior therapy, cognitive therapy, and medication in the treatment of major depressive disorder: a randomized clinical trial, posttreatment outcomes, and six-month follow-up.* *J Clin Psychol* 2008;64:728-46.
- Depkes. (2007). *Pharmaceutical Care* Untuk Penderita Gangguan Depresif, (online). Diakses di : <http://www.binfar.depkes.go.id>. (diakses tanggal 21 April 2020).
- Depbos. (2012). Depresi Penyebab Utama Gangguan Jiwa. Diakses di : <http://www.rehsos.depsos.go.id>. (diakses tanggal 21 April 2020).
- DeRubeis RJ, Hollon SD, Amsterdam JD, dkk. *Cognitive therapy vs medications in the treatment of moderate to severe depression.* *Arch Gen Psychiatry* 2005;62:409-16.
- Dimidjian S, Hollon SD, Dobson KS, dkk. *Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression.* *J Consult Clin Psychol* 2006;74:658-70.
- DiPiro J.T., Wells B.G., Schwinghammer T.L. and DiPiro C. V. (2015), *Pharmacotherapy Handbook*, Ninth Edit., McGraw-Hill Education Companies, Inggris.
- Driessens E, Hollon S. *Cognitive Behavioral Therapy for Mood Disorders: Efficacy, Moderators and Mediators.* Psychiatr Clin North Am. 2010 ; 33(3): 537–55.
- Epigee. (2009). *CBT for Post Traumatic Stress Disorder.* (online), diakses di : <http://www.epigee.org/ptsd-cbt.html> , (diakses tanggal 03 Mei 2020).
- Gartlehner G, Hansen RA, Morgan LC. (2011). *Comparative benefits and harms of second-generation antidepressants for treating major*

- depressive disorder: an updated meta-analysis. Ann Intern Med* 155:772-85.
- Gartlehner G, Thieda P, Hansen RA. (2008). *Comparative risk for harms of second-generation antidepressants: a systematic review and meta-analysis. Drug Saf*;31:851-65.
- Gosch, E. A., Flannery-Schroeder, E., Mauro, C. F., & Compton, S. N. (2006). *Principles of Cognitive-Behavioral Therapy for Anxiety Disorders in Children. Journal of Cognitive Psychotherapy*, 20(3), 247–262. <https://doi.org/10.1891/jcop.20.3.247>.
- Hawari D. (2001). Manajemen stres, cemas dan depresi. Jakarta: Balai Penerbit Fakultas Kedokteran Universitas Indonesia.
- Hegerl U, Hautzinger M, Mergl R, dkk. *Effects of pharmacotherapy and psychotherapy in depressed primary-care patients: a randomized, controlled trial including a patients' choice arm. Int J Neuropsychopharmacol* 2010;13:31-44.
- Institute of Medicine. (2001). *Crossing the quality chasm: a new health system for the 21st century*. National Academies Press.
- Ismail, R. I. dan Siste, K.(2010). Gangguan Depresi, Dalam Elvira,Silvia D., Hadisukanto, Gitayanti, Buku Ajar Psikiatri, Fakultas Kedokteran. Universitas Indonesia, Jakarta.
- Kando, J.C., Welss, B.G., and Hayes, P.E. (2005). *Pharmacoterapi A Pathophysiology Approach: Depressive Dissorder* 6th Ed Vol.2, 1235-1253.
- Kaplan dan Sadock. (2010). Buku ajar psikiatri klinis (diterjemahkan oleh Muttaqin H, Sihombing RNE). Edisi ke-2. Jakarta; Penerbit EGC
- Kaplan HI, Saddock BJ. Gangguan mood. Dalam: Sinopsis Psikiatri. Jilid 1 (Edisi Ketujuh). Jakarta: Binarupa Aksara (Bahasa Indonesia), 2010
- Kaplan, H. I dan Sadock, B. J. (1997). Ilmu Kedokteran Jiwa Darurat. Jakarta : Buku Kedokteran EGC.
- Karlsson, H., J. Hirvonen, J. Kajander, J. Markkula, H. Rasi-Hakala, J.K. Salminen,K. Nagren, S. Aalto, & J. Hietala. (2010). *Research letter: psychotherapy increases brain serotonin 5-HT1A receptors in patients with major depressive disorder. Psychological Medicine*, 40: 523–528.
- Katona, C., Cooper, C., Robrtson, M. (2012). At Glance Psikiatri, 2 nd , Jakarta: Erlangga, p. 18-19.

- Keck, P. E. (2005). *Bipolar Depression: A New Role For Atypical Antipsikotics.* <http://www.ncbi.nlm.nih.gov/pubmed/14700011>.
- Kennedy SH, Konarski JZ, Segal ZV, dkk. *Differences in brain glucose metabolism between responders to cognitive behavioral therapy and venlafaxine in a 16-week randomized controlled trial.* Am J Psychiatry 2007;164:778-88.
- Kennedy, S.H. dan Lam, R.W. (2003). *Enhancing Outcomes in The Management Resistant Depression: A Focus on Atypical Antipsikotic.* <http://www.ncbi.nlm.nih.gov/pubmed/15948765>.
- Lam RW, Parikh SV, Ramasubbu R, dkk. *Effects of combined pharmacotherapy and psychotherapy for improving work functioning in major depressive disorder.* Br J Psychiatry 2013;203:358-65.
- Lepine, Jean Pierre, Mike Briley, (2011). *The increase burden of depression Neuropsychiatr Dis trear.*(Suppl 1):S3-7.
- Mann, J. J. (2005). *The Medical Management of Depressi, The New England Journal of Medicine*, number 17, volume 353: 1819 – 1834.
- Martin, P.F. (2010). CBT 27 Maret 2010. Diakses di : <http://www.minddisorders.com/Br-Del/Cognitive-behavioral-therapy.html>. (diakses tanggal 03 Mei 2020).
- Maulida, A. (2012). Gambaran Tingkat Depresi Pada Mahasiswa Program Sarjana Yang Melakukan Konseling Di Badan Konseling Universitas Indonesia. Skripsi. Jakarta: Universitas Indonesia.
- Mayasari,W.T. (2013). “Gambaran Umum Depresi”. Denpasar. Diambil dari <https://www.google.com/search?q=gambaran+umum+depresi+mayasari&i&oq=gambaran+umum+depresi+mayasari&aqs=chrome..69i57.13013j0j4&sourceid=chrome&ie=UTF-8>. (diakses tanggal 02 Mei 2020).
- Miranda J, Chung JY, Green BL, dkk. *Treating depression in predominantly low-income young minority women: a randomized controlled trial.* JAMA 2003;290:57-65.
- Mojtabai R, Olfson M. (2014). *National trends in long-term use of antidepressant medications: results from the U.S. National Health and Nutrition Examination Survey.* J Clin Psychiatry ;75:169-77. <https://www.ncbi.nlm.nih.gov/pubmed/24345349>.
- Mynors-Wallis LM, Gath DH, Day A, dkk. *Randomised controlled trial of problem solving treatment, antidepressant medication, and combined treatment for major depression in primary care.* BMJ 2000;32:26-30.

- Neal, M. J. (2006). *At Glane Farmakologi Medis*, Edisi kelima, diterjemahkan oleh surapari, J., dr., & Safitri, A., Gelora Aksara Pratama, Jakarta.
- O'Hea, E., Houseman, J., Bedek, K., & Sposato, R. (2009). *The use of cognitive behavioral therapy in the treatment of depression for individuals with CHF*. *Heart Failure Reviews*, 14(1), 13–20.
- Oemarjoedi, A. K. (2003). Pendekatan *cognitive behavior* dalam psikoterapi. *Jakarta: Kreativ Media*.
- Pilkington K.(2010). *Anxiety, depression and acupuncture: a review of the clinical research*. *Autonomic neuroscience: basic and clinical research*. *Autonomic neuroscience* ;157:91-5.
- Prayitno. (2008). *Farmakologi Dasar & Klinik*, Lilian Batubara (eds), 129 – 130 Penerbit Lenskopi, Jakarta.
- Riset Kesehatan Dasar (Risksdas). (2018). Badan Penelitian dan Pengembangan Kesehatan Kementerian RI tahun 2018. Diakses di : http://kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-risksdas-2018_1274.pdf, (diakses tanggal 25 April 2020).
- Sadock BJ, Sadock VA. Kaplan dan Sadock's (2007). *synopsis of psychiatry behvioral sciences/clinical psychiatry*. 10th ed. Philadelphia: Lippincott Williams & Wilkins.
- Santrock, J. W. (2003). Adolescence: Perkembangan Remaja (alih bahasa Shinto B. Adelar dan Sherly Saragih). *Jakarta: Erlangga*.
- Segal ZV, Kennedy S, Gemar M, dkk. *Cognitive reactivity to sad mood provocation and the prediction of depressive relapse*. *Arch Gen Psychiatry* 2006;63:749-55.
- Shamsaei F, Rahimi A, Zarabian MK, dkk. *Efficacy of pharmacotherapy and cognitive therapy, alone and in combination in major depressive disorder*. *Hong Kong J Psychiatr* 2008;18:76-80.
- Stuart,G.W. (2009). *Principles and practice of psychiatric Nursing*. 9th ed. Missouri: Mosby, inc.
- Sukandar, E. Y., Andrajati, R., Sigit, I. J., Adnyana, K. I., Setiadi, P. A. A., Kusnandar. (2009). *ISO Farmakoterapi*, Cetakan kedua, PT. ISFI Penerbitan, Jakarta.
- Teter, C. S., Kando, J. C., Wells, B. G., dan Hayes, P. E. (2007). *Depressive Disorder*,dalam Dipiro, J. T., Talbert, R. L., Yee, G. C., Matzke, G. R., Wells, B. G.,& Posey Micheal, L., (eds), *Pharmacotherapy A*

Pathophysiologic Approach,7th Edition, Appleton and lange, New York.

Tjay, T.H dan Rahardja, K. (2007). Obat-Obat Penting Khasiat, Penggunaan dan Efek-Efek Sampingya, PT. Elex Media Komputindo, Jakarta.

Townsend, A. K., Clark, A. B., McGowan, K. J., Buckles, E. L., Miller, A. D., dan Lovette, I. J. (2009). *Disease-mediated inbreeding depression in a large, open population of cooperative crows*. *Proceedings of the Royal Society B: Biological Sciences*, 276(1664), 2057-2064.

WHO. (2020). *Depression*. diakses di : <https://www.who.int/news-room/fact-sheets/detail/depression> (diakses tanggal 02 Mei 2020).

Xing S, Dipaula BA, Lee HY. (2011). *Failure to fill electronically prescribed antidepressant medications: a retrospective study*. *Prim Care Companion CNS Disord* ;13:pii: PCC.10m00998.

Yatham, L. N., 2004, *Newer Antikonvulsants in The Treatment of Bipolar Disorder*. <http://www.ncbi.nlm.gov/pubmed/1524330>.

Yuniastuti. (2013). *Evaluasi Terapi Obat Antidepresan Pada Pasien Depresi di RS Jiwa Dr. Soejarwadi Provinsi Jawa Tengah tahun 2013*.